Note on Legendre decomposition of the Pontryagin density in Kerr

Leo C. Stein*
Center for Radiophysics and Space Research, Cornell University, Ithaca, NY 14853 USA
(Dated: July 4, 2014)

In [1] ("Scalar field excited around a rapidly rotating black hole in Chern-Simons modified gravity"), Konno and Takahashi have recently developed some analytical results for the scalar field about a Kerr black hole in the decoupling limit of dynamical Chern-Simons gravity. This involved a decomposition of the source (the Pontryagin density) in terms of Legendre polynomials. Here we give a two-line expression for this decomposition which simplifies their quadruple sum. Our expressions are rational polynomials multiplying Legendre functions of the second kind, or equivalently rational polynomials multiplying hypergeometric functions.

PACS numbers: 04.50.Kd,02.30.Gp

arXiv:1407.0744v1 [gr-qc] 2 Jul 2014

In [1], Konno and Takahashi (henceforth KT) attempted to find an analytical solution for the scalar field ϑ which satisfies

$$\Box\vartheta \propto {}^*RR \tag{1}$$

where ${}^*RR = -{}^*R^{abcd}R_{abcd} = -\frac{1}{2}\epsilon^{abef}R_{ef}{}^{cd}R_{abcd}$ is the Pontryagin density. This is one of the equations of motion of dynamical Chern-Simons (dCS) gravity [2, 3], the other being the modified Einstein equation, which also includes corrections from ϑ. In the decoupling limit about some background (GR) solution, the wave operator of Eq. (1) becomes the background wave operator and the source term is evaluated on the background spacetime. When seeking rapidly-rotating black hole solutions we use the Kerr spacetime [4, 5] as the background solution, and impose stationarity and axisymmetry on ϑ. Then we desire to solve

$$\left[\frac{\partial}{\partial r}\left(\Delta\frac{\partial}{\partial r}\right) + \frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\left(\sin\theta\frac{\partial}{\partial\theta}\right)\right]\vartheta = C\Sigma{}^*RR\,, \tag{2}$$

where $\Delta \equiv r^2 + a^2 - 2Mr$, $\Sigma \equiv r^2 + a^2\cos^2\theta$, C is some constant related to the coupling strength of dCS, and in Kerr we have

$${}^*RR = \frac{96M^2acr}{\Sigma^6}(3r^2 - a^2c^2)(r^2 - 3a^2c^2)\,, \tag{3}$$

where $c = \cos\theta$.

The differential operator on the LHS of Eq. (2) separates, and the eigenfunctions of the homogeneous angular equation are simply the Legendre polynomials $P_l(c)$. Therefore we consider the decomposition

$$S \equiv C\Sigma{}^*RR = \sum_{l=0}^{\infty} S_l(r)P_l(\cos\theta) \tag{4}$$

where

$$S_l(r) = \frac{2l+1}{2}\int_{-1}^{+1} S(r,c)P_l(c)dc\,. \tag{5}$$

The source is odd in c, so we will only need to consider odd l.

For simplicity we define

$$S_l(r) = 96CM^2\frac{2l+1}{2}I_l(r) \tag{6}$$

$$I_l(r) \equiv \int_{-1}^{+1}\frac{acr(3r^2 - a^2c^2)(r^2 - 3a^2c^2)}{(r^2 + a^2c^2)^5}P_l(c)dc\,, \tag{7}$$

which appears as a source term for the separated inhomogeneous radial equation for $\vartheta_l(r)$, defined similarly to $S_l(r)$.

I. TWO-LINE EXPRESSION FOR $I_l(r)$

KT give their $S_{2n+1}(\tilde{r})$ (not quite equal to ours, see Sec. II) in their Eqs. (29-36) in terms of a quadruple sum of rational polynomials and the arctan function. We find two-line expressions for $I_l(r)$ in terms of special functions:

$$\begin{aligned} I_l(r) = &-\frac{(l+1)}{24(a^2+r^2)^4}\left\{\frac{r}{a}\left[a^4(l^3 - l^2 - 16l + 4) + 2a^2r^2\left(l^2(l+3) - 16\right) + r^4(l+2)^2(l+3)\right]\left[Q_l\left(+\tfrac{ir}{a}\right) + Q_l\left(-\tfrac{ir}{a}\right)\right]\right. \\ &\left. + 2\left[a^4\left(l^2 + l - 2\right) - 2a^2r^2\left(l^2 + l - 8\right) - 3r^4\left(l^2 + l + 2\right)\right]\frac{1}{i}\left[Q_{l+1}\left(+\tfrac{ir}{a}\right) - Q_{l+1}\left(-\tfrac{ir}{a}\right)\right]\right\}, \end{aligned} \tag{8}$$

* Einstein fellow; leostein@astro.cornell.edu

or equivalently,

$$I_l(r) = \frac{\varepsilon(l)\sqrt{\pi}\Gamma(l+2)\,(a/r)^l}{3\,(2)^{l+3}\,(a^2+r^2)^4\,\Gamma(l+\frac{3}{2})}\Big\{\left[a^4(l^3-l^2-16l+4)+2a^2r^2\left(l^2(l+3)-16\right)+r^4(l+2)^2(l+3)\right]F(\tfrac{l+1}{2},\tfrac{l+2}{2};l+\tfrac{3}{2};-\tfrac{a^2}{r^2}) \\ -\frac{2(l+1)a^2}{(2l+3)r^2}\left[a^4\left(l^2+l-2\right)-2a^2r^2\left(l^2+l-8\right)-3r^4\left(l^2+l+2\right)\right]F(\tfrac{l+2}{2},\tfrac{l+3}{2};l+\tfrac{5}{2};-\tfrac{a^2}{r^2})\Big\} \tag{9}$$

where $\varepsilon(l) = +1$ for $l = 1, 5, 9, \ldots$, $\varepsilon(l) = -1$ for $l = 3, 7, 11, \ldots$, and $\varepsilon(l) = 0$ for even l. Below we describe how to find these expressions.

We make use of the identity[1]

$$\int_{-1}^{+1} c^k(z-c)^{-1}(1-c^2)^{m/2}P_l^m(c)dc \\ = (+2)(z^2-1)^{m/2}Q_l^m(z)z^k \tag{10}$$

where $m \le l, k = 0, 1, \ldots l-m$, and z is in the complex plane with a cut along $(-1, +1)$ on the real axis. To use this identity, we will have $m = 0 = k$,

$$\int_{-1}^{+1}\frac{P_l(c)}{z-c}dc = +2Q_l(z) \tag{11}$$

with the same restriction on z as above. To get Eq. (7) into a form where Eq. (11) may be applied, use a complex partial fractions decomposition for the rational polynomial (i.e. the denominator $(r^2+a^2c^2)^5$ is an irreducible polynomial over $\mathbb{R}$ but it is reducible over $\mathbb{C}$). The decomposition is

$$\frac{acr(3r^2-a^2c^2)(r^2-3a^2c^2)}{(r^2+a^2c^2)^5} = \frac{r}{2(ac-ir)^5}+\frac{r}{2(ac+ir)^5} \\ -\frac{i}{4(ac-ir)^4}+\frac{i}{4(ac+ir)^4}\,. \tag{12}$$

Now the original integral I_l has been converted to two integrals of the form $\int P_l(c)/(ac \pm ir)^n dc$ where $n = 4, 5$. The power n may be reduced through integration by parts, i.e. integrating $(ac \pm ir)^{-n}dc$ while differentiating $P_l(c)$. After again performing a partial fractions decomposition, this creates two types of terms. First, terms of the form $\int P_{l'}(c)/(ac \pm ir)^{n'}dc$ where $n' = 1, 2, \ldots, n-1$ and $l' = l, l+1$, via [8]

$$(1-x^2)\frac{dP_\nu^\mu(x)}{dx} = (\mu-\nu-1)P_{\nu+1}^\mu(x)+(\nu+1)xP_\nu^\mu(x)\,. \tag{13}$$

Second, terms of the form $\int P_{l'}(c)/(c \pm 1)dc$. The former terms with $n' = 1$ may be evaluated directly with Eq. (11) and the other n' may be repeatedly integrated by parts as just described. The remaining terms of the form $\int P_{l'}(c)/(c \pm 1)dc$ can all be combined together into integrals of the form

$$\int\frac{P_{l'-1}(c)-P_{l'+1}(c)}{1-c^2}dc\,. \tag{14}$$

Here the integrand is subject to the identity

$$P_{n-1}(x)-P_{n+1}(x) = \frac{(2n+1)\left(1-x^2\right)P_n'}{n(n+1)} \tag{15}$$

which immediately yields

$$\int_a^b\frac{P_{l'-1}(c)-P_{l'+1}(c)}{1-c^2}dc = \frac{(2n+1)}{n(n+1)}\left[P_n(b)-P_n(a)\right]\,. \tag{16}$$

Applying these identities allows us to integrate $I_l(r)$ and gives Eq. (8).

Though the argument of Q_l is purely imaginary in Eq. (8), the combinations $(Q_l(ix)+Q_l(-ix))$ and $\frac{1}{i}(Q_l(+ix)-Q_l(-ix))$ are purely real. This can be seen with the identity [8]

$$Q_l(z) = \frac{\sqrt{\pi}\Gamma(l+1)}{(2z)^{l+1}\Gamma(l+\frac{3}{2})}F\left(\frac{l+1}{2},\frac{l+2}{2};l+\frac{3}{2};\frac{1}{z^2}\right)\,. \tag{17}$$

Using this identity gives Eq. (9) which is manifestly real.

II. CHECKS AND COMPARISONS

For any given l, typical computer algebra systems (such as MATHEMATICA) can perform the explicit integral $I_l(r)$, since it is nothing but a rational polynomial function. We have checked that Eq. (9) agrees with the explicit evaluation of these integrals for a large number of l's.

We have also compared our expressions (given in Appendix A) with those given in KT. We have verified the relationship

$$S_l^{\rm KT} = \frac{2l+1}{2}I_l \tag{18}$$

where $S_l^{\rm KT}$ are the expressions given in Appendix A of KT. This suggests that KT have dropped the factor of $96C$ (they scale all dimensional quantities by M). Their expressions should be multiplied by this factor, which they take as $3/2\pi$.

[1] This identity is correct in the fourth edition [6] but incorrect in the seventh edition [7]. I did not have access to other editions to check where the error was made.

ACKNOWLEDGMENTS

The author would like to acknowledge Barry Wardell for helpful discussions. LCS acknowledges that support for this work was provided by the National Aeronautics and Space Administration through Einstein Postdoctoral Fellowship Award Number PF2-130101 issued by the Chandra X-ray Observatory Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of the National Aeronautics Space Administration under contract NAS8-03060.

Appendix A: Source moments for small l

$$I_1 = +2(a^2+r^2)^{-4} r a (r^2-a^2) \tag{A1}$$

$$I_3 = -\tfrac{1}{6}(a^2+r^2)^{-4} r a^{-3}\left[57a^6+73a^4r^2 + 55a^2r^4+15r^6\right] + \tfrac{5}{2}a^{-4}\arctan\left(\tfrac{a}{r}\right) \tag{A2}$$

$$I_5 = +\tfrac{1}{12}(a^2+r^2)^{-4} r a^{-5}\left[1047a^8+5533a^6r^2 + 9583a^4r^4+7035a^2r^6+1890r^8\right] - \tfrac{35}{4}a^{-6}(a^2+18r^2)\arctan\left(\tfrac{a}{r}\right) \tag{A3}$$

$$I_7 = -\tfrac{1}{16}(a^2+r^2)^{-4} r a^{-7}\left[4805a^{10}+56665a^8r^2 + 188298a^6r^4+270354a^4r^6 + 179025a^2r^8+45045r^{10}\right] + \tfrac{315}{16}a^{-8}(a^4+44a^2r^2+143r^4)\arctan\left(\tfrac{a}{r}\right) \tag{A4}$$

[1] K. Konno and R. Takahashi, (2014), arXiv:1406.0957 [gr-qc].
[2] R. Jackiw and S. Y. Pi, Phys.Rev. **D68**, 104012 (2003), arXiv:gr-qc/0308071 [gr-qc].
[3] S. Alexander and N. Yunes, Phys.Rept. **480**, 1 (2009), arXiv:0907.2562 [hep-th].
[4] R. P. Kerr, Phys.Rev.Lett. **11**, 237 (1963).
[5] M. Visser, (2007), arXiv:0706.0622 [gr-qc].
[6] I. S. Gradshteyn and I. M. Ryzhik, *New York: Academic Press, 1965, 4th ed., edited by Geronimus, Yu.V. (4th ed.); Tseytlin, M.Yu. (4th ed.)* (1965).
[7] I. S. Gradshteyn, I. M. Ryzhik, A. Jeffrey, and D. Zwillinger, *Table of Integrals, Series, and Products, Seventh Edition by I. S. Gradshteyn, I. M. Ryzhik, Alan Jeffrey, and Daniel Zwillinger. Elsevier Academic Press, 2007. ISBN 012-373637-4* (2007).
[8] DLMF, "NIST Digital Library of Mathematical Functions," http://dlı Release 1.0.6 of 2013-05-06, online companion to [9].
[9] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, eds., *NIST Handbook of Mathematical Functions* (Cambridge University Press, New York, NY, 2010) print companion to [8].